



Hy Laboratories Ltd.

6 Menachem Plaut St., Park Tamar, Rehovot 7670606, Israel

Tel. +972.8.9366475 Fax. +972.8.9366474

hylabs@hylabs.co.il | www.hylabs.co.il

**Cat No. CD013 - SDA+LECITHIN+POLYSORBATE 80  
(IRRADIATED)****Product Type:** CONTACT DISHES**Application-** culture medium used in microbiology for the isolation and cultivation of fungi, especially molds and yeasts. Each component in the medium serves a specific.**Intended Use** – Sabouraud Dextrose Agar is a general-purpose medium used in the cultivation of yeasts, molds and aciduric microorganisms. Cultivation of pathogenic and nonpathogenic fungi, particularly dermatophytes. SDA+LECITHIN+POLYSORBATE 80 provides a nutrient-rich and selective environment for the cultivation of fungi, including molds and yeasts. The medium supports the growth of a wide range of fungi while inhibiting the growth of most bacteria due to its acidic pH. Lecithin and Polysorbate 80 may be added to enhance the recovery and growth of fungi with specific nutritional requirements, especially those that thrive on lipid substrates. Fungal cultures on this medium are typically incubated at appropriate temperatures for the isolation and identification of fungal species. The resulting fungal colonies can be examined, studied, and identified for various research, clinical, and diagnostic purpose**Composition g/L**

Peptic digest of Animal Tissue 5g/L,

Pancreatic digest of Casein 5g/L,

Dextrose 40g/L,

Agar 15g/L

LECITHIN – 0.7g/L

TWEEN 80 - 5 ml/L

**Storage:** 2-25°C**Appearance:** light amber**pH Range:** 5.4 - 5.8**Gamma Irradiation (kGy) Range:** 7 - 15**Warning and Precautions** - For professional use only. Follow good microbiological lab practices while handling specimens and culture. Do not use Petri dishes if they show evidence of microbial contamination, discoloration, drying, cracking, or other signs of deterioration. Avoid freezing and overheating. The Petri Dishes may be used / inoculated up to the expiration date and incubated for the recommended incubation times. After use and prior to discarding, specimen containers and all contaminated material, including the used culture media and contaminated culture containers, must be sterilized or incinerated by validated procedures. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.



Hy Laboratories Ltd.

6 Menachem Plaut St., Park Tamar, Rehovot 7670606, Israel

Tel. +972.8.9366475 Fax. +972.8.9366474

hylabs@hylabs.co.il | www.hylabs.co.il

## Performance Testing Results

Spreading from estimated inoculum 10-100 cfu.

TEST	ATCC	Incubation Temp.(°C)	Incubation Cond.	Reaction 1	Recovery (%)
<i>Candida albicans</i>	10231	20-25 °C	Aerobic, 3-5 days	Pass	70-200
<i>Penicillium expansum</i>	7861	20-25 °C	Aerobic, 3-5 days	Pass	70-200
<i>Aspergillus brasiliensis</i>	16404	20-25 °C	Aerobic, 3-5 days	Pass	70-200

## Neutralizer's Effectiveness Test:

Disinfectant: Glyoxal 1.2% glutardialdehyde 0.05% didecyl-dimethyl-ammonium chloride 0.75%

TEST	ATCC	Incubation Temp.(°C)	Incubation Cond.	Reaction
<i>Candida albicans</i>	10231	20-25 °C	Aerobic, 3-5 days	Confluent growth
<i>Penicillium expansum</i>	7861	20-25 °C	Aerobic, 3-5 days	Confluent growth
<i>Aspergillus brasiliensis</i>	16404	20-25 °C	Aerobic, 3-5 days	Confluent growth